

Factors associated with electronic cigarettes use among adolescents in Jakarta, Indonesia

by Ridhwan Fauzi

Submission date: 28-Mar-2023 05:28AM (UTC-0400)

Submission ID: 2048917679

File name: JHR-Ecig.pdf (120.7K)

Word count: 5178

Character count: 28732

Factors associated with electronic cigarettes use among adolescents in Jakarta, Indonesia

Ridhwan Fauzi and Chitlada Areesantichai

College of Public Health Sciences, Chulalongkorn University, Bangkok, Thailand

Abstract

Purpose – This study aimed to examine factors associated with electronic cigarette use among adolescents aged 15–19 in Jakarta, Indonesia.

Design/methodology/proach – The study was a school-based survey involving 1,318 students from 14 high schools in Jakarta. A multistage cluster sampling methodology was used. The authors used the chi-square test and multiple logistic regression to examine the association between electronic cigarette use and sociodemographic, social influences, health risk perceptions, availability, affordability and conventional cigarette smoking status.

Findings – Overall, 6.3% of females and 29% of males reported ever having used electronic cigarettes. Electronic cigarette use was independently associated with sex, school locations, conventional cigarette smoking status, peer use, availability and perceptions that electronic cigarettes aid conventional cigarette smoking cessation. Compared with non-smokers, lifetime (AOR: 8.740, 95% CI: 5.126–14.901) and current conventional cigarette smokers (AOR: 18.380, 95% CI: 10.577–31.938) were more likely to use electronic cigarettes.

Social implications – The use of electronic cigarettes among adolescents was considerably high in this study. Therefore, the tobacco control policy should be extended to regulate the marketing and use of all types of tobacco products and not just conventional cigarettes.

Originality/value – The study explored beyond individual lifestyle factors that contributed to electronic cigarette use in Indonesia such as affordability, school locations and availability of electronic cigarettes. These issues have not been specifically discussed in previous studies.

Keywords Adolescents, Electronic cigarettes, Tobacco products, Vaping, Smoking cessation, Indonesia

Paper type Research paper

Introduction

The electronic nicotine delivery system (ENDS) or electronic cigarette is a major public health issue in many parts of the world [1]. The prevalence of electronic cigarette smoking varies across the WHO member countries ranging from 0.7% to 23.4% [2]. Nevertheless, studies find that electronic cigarette smoking has been increasing rapidly in recent years [3,4]. Basic health research 2018 reported that the prevalence of current electronic cigarette use is 2.8%, which consists of 2.8% in males and 2.7% in females [5]. The prevalence may still be lower than the US or European countries, but the government should be aware of the vaping trend, particularly among youth.

Electronic cigarettes have raised the attention of medical and public health professionals due to its potential benefit and harm [6]. The industry passionately advertises electronic cigarettes as an aid to conventional cigarette cessation [7]. However, the evidence is still

© Ridhwan Fauzi and Chitlada Areesantichai. Published in *Journal of Health Research*. Published by Emerald Publishing Limited. This article is published under the Creative Commons Attribution (CC BY 4.0) licence. Anyone may reproduce, distribute, translate and create derivative works of this article (for both commercial and non-commercial purposes), subject to full attribution to the original publication and authors. The full terms of this licence may be seen at <http://creativecommons.org/licences/by/4.0/legalcode>

This study was partially supported by the 90th Anniversary of Chulalongkorn University Scholarship.



30
3 rking regarding the effectiveness of electronic cigarettes as a cessation tool [8]. In addition, the safety and long-term health effects of electronic cigarettes used are not yet apparent [9,10]. Prior studies reported that electronic cigarette aerosols contain similar hazardous components as found in conventional cigarettes such as nickel, chromium, cadmium, tin, carbonyl compounds, volatile organic compounds and aluminum [11–13]. A study in Surabaya, a city in Indonesia, found high formaldehyde concentrations in electronic cigarette users' vapors [14]. This chemical compound is carcinogenous and leads to a fatal health condition in the long-term [11]. Besides, the particulate meter (PM2.5) concentration in electronic cigarette user's homes were higher compared to smoke-free homes [15]. Therefore, the emission from electronic cigarettes has a potentially adverse effect on second-hand smokers.

Electronic cigarette industries expend a wide variety of marketing channels to promote their products including magazines and print media ads, online marketplaces, points of sales ads, television commercials, sponsorship and social media [11]. The marketing techniques and themes are similar to those that have been found in conventional cigarettes [11]. In Indonesia, electronic cigarettes are mainly traded in the online platform (35.3%) and vape stores (64.7%) [16]. Vape shops and electronic cigarette ambassadors also use Instagram, the most popular visual social media platform in Indonesia, that may appeal to the youth [17]. A recent study revealed that Indonesia shared the second-largest Instagram post on e-liquid marketing of any country [18].

11
The evidence showed that the demand for electronic cigarettes is excessively sensitive to the owned price change [19]. The price of electronic cigarettes may be substantially higher than conventional cigarettes in Indonesia. For instance, a pack of A Mild, the most popular cigarette brand, cost around 1.6 USD, while a device kit of a popular electronic cigarette brand sold on Blibli.com was approximately 12.65 USD [20,21]. The electronic cigarettes tax and pricing policies are also not well regulated [16]. Thus, the industry is able to develop a pricing and promotional strategy to make the product more affordable and accessible to the youth.

Indonesian Food and Drug Administration (FDA) has declared electronic cigarettes as a dangerous product [17]. Nevertheless, the trend of electronic cigarette use among adolescents continues to steadily increase in Indonesia [22]. In addition, the multinational electronic cigarette industry has expanded quite prominently into Indonesia's market recently [16]. Until now, there was no specific regulation about electronic cigarette marketing and use [17]. Overall, the literature on electronic cigarettes remains scarce in Indonesia. Therefore, this study aimed to investigate factors associated with electronic cigarette use among adolescents aged 15–19 years old in Jakarta. The findings would provide a framework to develop an effective intervention program to prevent youth from the damaging uptake of electronic cigarettes.

Materials and methods

The study was a school-based survey conducted between April and May 2015. The study area was Jakarta Province, the capital city of Indonesia. There are approximately 377,216 students from 1,263 high schools across Jakarta Province [23]. The minimum sample size was 768 students determined by sample size formula to estimate proportion with 0.05 level of significance and a 0.05 acceptable error. We only included general public schools as our study sites. The study used a multistage cluster sample of high school students in Jakarta. First, we randomly selected eight schools from the suburban area and six schools from the downtown area. Second, we randomly selected three classes from grades ten and eleven from each school. Finally, 25 students from the selected classes were recruited as study participants. The researcher explained the study objectives and informed consent process before data collection. The students who were willing to participate were required to sign a consent form.

The data were obtained by a self-administered questionnaire that took approximately 15 min to complete.

Three tobacco control experts from Thailand assessed the validity of the questionnaire. The total item objective congruence (IOC) score was 0.858. Sociodemographic characteristics of study participants included age, sex, school location (suburban vs ³⁵ wntown) and parents' education attainment (primary vs secondary vs higher). Electronic cigarette smoking status was measured by asking two questions: (1) "Have you ever tried or experimented with smoking an electronic cigarette, even one or two puffs?" (2) "During the past 30 days, did you smoke elect¹⁴ c cigarettes?". The responses included "no" and "yes" options. Study participants who answered "yes" to both questions were defined as current smokers. We measured the social influences by inquiring about the electronic cigarette smoking behaviors of family members and peers.

The health risk ¹² eptions were measured by five positively framed questions; (1) electronic cigarettes help cut down the number of cigarettes smoked; (2) aids in quitting cigarette smoking; (3) doesn't contain toxic chemicals that have been found in combustible cigarettes; (4) less harmful and (5) less addictive. Three ¹³ ons were used to measure health risk perception namely "yes", "no", and "don't know". The Cronbach's alpha score for the health risk perception variable was 0.794. Conventional cigarette-related questions were "Have you ever tried or experimented with cigarette smoking, even one or two puffs?" (yes/no) and "During the past 30 days, did you smoke a cigarette?" (yes/no). For assessing conventional cigarette smoking status, two questions were asked: (1) "Have you ever tried or experimented with conventional cigarette smoking, even one or two puffs?" and (2) "During ¹⁵ past 30 days, did you smoke conventional cigarettes?". The participants who had smoked one or more cigarettes before but not in the past 30 days were considered to be lifetime smokers. The last part of the questionnaire measured participants' perceptions about the availability and affordability of electronic cigarettes. The availability was explored by participants' perception of difficulty getting electronic cigarettes. The affordability was assessed by inquiring whether electronic cigarette prices were affordable according to the stu²³ participant's perspective.

Descriptive statistics were used to summarize the general characteristics of the participants. Electronic cigarette use variables were transformed into binary classification, namely users and non-users. Users of electronic cigarettes consisted of historic lifetime and current smokers. A chi-square test was performed for bivariate analysis. Independent variables' significance at $p < 0.25$ in bivariate analysis was included in the multivariate model. Multiple logistic regressions estimated an adjusted odds ratio (AOR) and 9¹⁷ confidence interval (CI). Statistical significance was considered at p -value < 0.05 . All statistical analyses were performed by SPSS statistics version 22.0 (IBM Corp, Armonk, NY ²⁴ SA).

The protocol was approved by the Institutional Review Board, ¹⁹ Atma Jaya Catholic University of Indonesia, number 404/III/LPPM-PM.10. 05/04/2015.

Results ²⁷

Table 1 shows the general characteristics of the study participants by sex. Of the 1,318 participants, about 62.8% were female. The average age was 16 years old among both males and females (16.08 ± 0.71 vs 16.21 ± 0.72). The current conventional cigarette smokers were approximately 8.5% female and 29.2% male. We estimated that ³⁶ current electronic cigarette smokers among females and males were 0.6% and 8.2%. The rate of the current use of electronic cigarettes of the family members was low among both males and females at 3.1% and 2.8%. About one-fifth (20%) of the males had at least one close friend who currently smoked electronic cigarettes. A total of 38% of the participants reported that they easily

Characteristics <i>n</i> (%) or mean \pm SD	Female <i>n</i> = 828	Male <i>n</i> = 490
Age	16.08 \pm 0.71	16.21 \pm 0.72
<i>School location</i>		
Suburban	491 (59.3)	287 (58.6)
Downtown	337 (40.7)	203 (41.4)
<i>Parents' education</i>		
Completed primary or less	33 (4.0)	12 (2.4)
Secondary	437 (52.8)	234 (47.8)
Higher	358 (43.2)	244 (49.8)
<i>Conventional cigarettes smoking</i>		
None	643 (77.7)	179 (36.5)
Lifetime	115 (13.9)	168 (34.4)
Current	70 (8.5)	143 (29.2)
<i>Electronic cigarettes smoking</i>		
None	776 (93.7)	348 (71.0)
Lifetime	47 (5.7)	102 (20.8)
Current	5 (0.6)	40 (8.2)
<i>Family use</i>		
No	805 (97.2)	475 (96.9)
Yes	23 (2.8)	15 (3.1)
<i>Peer use</i>		
No	695 (83.9)	392 (80)
Yes	133 (16.1)	98 (20)
<i>Availability</i>		
Difficult	506 (61.1)	300 (61.2)
Easy	322 (38.9)	190 (38.8)
<i>Affordability</i>		
No	608 (73.4)	347 (70.8)
Yes	220 (26.6)	143 (29.2)

Table 1.
Characteristics of
study participants
by sex

found electronic cigarettes. More than a quarter (29.2%) of the males perceived that electronic cigarette prices were affordable.

A total of 27.3% of electronic cigarette users believed that electronic cigarettes were a smoking cessation aid, 30.1% reported they could help cut down the number of cigarettes smoked and 19.7% reported they were less addictive. More electronic cigarette users reported that electronic cigarettes were less harmful (20.2%) than non-users (9.1%). Health risk perceptions were significantly associated with ever having smoked electronic cigarettes at $p < 0.001$ (Table 2).

Table 3 indicates the bivariate analysis of ever having used electronic cigarettes. Of electronic cigarette users, 3% had never used conventional cigarettes. Age was the only variable that did not significantly associate with ever having used electronic cigarettes in the bivariate analysis. However, the p -value was still less than 0.25. Thereby, all variables were included in the multivariate analysis.

Table 4 presents multiple logistic regression results. Having ever used electronic cigarettes were independently associated with the male sex (AOR: 3.522, 95% CI: 2.316–5.757), downtown school locations (AOR: 1.711, 95% CI: 1.153–2.538), having peers who

currently ¹⁰ electronic cigarettes (AOR: 2.073, 95% CI: 1.311–3.277), easy access to get electronic cigarettes (AOR: 2.370, 95% CI: 1.526–3.679) and the perception that electronic cigarettes were an aid to conventional cigarettes cessation (AOR: 5.197, 95% CI: 2.099–13.106). ³ compared with non-smokers, both lifetime smokers (AOR: 8.740, 95% CI: 5.126–14.901) and current conventional cigarette smokers (AOR: 18.380, 95% CI: 10.577–31.938) were more likely to ever use electronic cigarettes.

6

Table 2.
³⁸th risk perceptions of electronic cigarettes by electronic cigarettes smoking status

Health risk perceptions of electronic cigarettes	Electronic cigarettes use status	
	User (n = 193) n (%)	Non-user (n = 1,125) n (%)
Help cut down number of cigarettes*	58 (30.1)	105 (9.3)
Aid quit cigarette smoking*	53 (27.5)	72 (6.4)
Does not contain toxic chemical*	30 (15.5)	82 (7.3)
Less harmful*	39 (20.2)	102 (9.1)
Less addictive*	38 (19.7)	104 (9.2)

Note(s): *p < 0.001

Table 3.
Bivariate analysis of ever using electronic cigarettes

Independent variables	Electronic cigarettes use		p-value
	User (n = 193) n (%)	Non-user (n = 1,125) n (%)	
<i>Age (years)</i>			0.05
15–16	70 (17.5)	329 (82.5)	
17–19	123 (13.4)	796 (86.6)	
<i>Sex</i>			<0.001
Male	142 (29.0)	348 (71.0)	
Female	52 (6.2)	777 (93.8)	
<i>School locations</i>			<0.001
Downtown	104 (19.3)	436 (80.7)	
Suburban	89 (11.4)	689 (88.6)	
<i>Parents' education</i>			0.002
≤ Primary	44 (97.8)	1 (2.2)	
Secondary	85 (12.7)	586 (87.3)	
College degree	107 (17.8)	495 (82.5)	
<i>Cigarette use</i>			<0.001
None	25 (3.0)	797 (97.0)	
Lifetime	74 (26.1)	209 (73.9)	
Current	94 (44.1)	119 (55.9)	
<i>Family use</i>			0.001
No	1,100 (85.9)	180 (14.1)	
Yes	13 (34.2)	25 (64.8)	
<i>Peer use</i>			<0.001
No	139 (12.8)	948 (87.2)	
Yes	54 (23.4)	177 (76.6)	
<i>Availability</i>			<0.001
Difficult	78 (9.7)	728 (90.3)	
Easy	115 (22.5)	397 (77.5)	
<i>Affordability</i>			<0.001
No	111 (11.6)	844 (88.4)	
Yes	82 (22.6)	281 (77.4)	

Discussion

The electronic cigarette smoking prevalence among male adolescents who had ever used them was about 29%, and 8.2% included current users according to this study. The findings are slightly higher than the rate reported from other studies in Poland (21.8% of users), Ontario (18.6% of users) and South Korea (7.8% current use) [24–26]. The fact is unsurprising because those areas have strong regulations on the restriction of electronic cigarette marketing and use [27]. Presently, there is no specific regulation in Indonesia on electronic cigarette control at the national and sub-national levels [17].

The Global Adult Tobacco Survey 2011 reported that the prevalence of electronic cigarettes is about 0.5% in males and less than 0.1% in females [28]. The rate markedly increased by more than ten times based on our study in 2015. The soaring prevalence of electronic cigarette smoking may also relate to the expansion of multinational companies to Indonesia [16,29]. Indonesia has poor tobacco control regulations compared to other ASEAN countries [21]. Besides, the country has a large young population that is a promising market for the tobacco industry [30]. China vape manufacturers pave the way for creating electronic cigarette demand in Indonesia [31]. Nevertheless, US and local products have also started to compete in the market [16,17,31]. Recently, the Minister of Industry announced that Phillip Morris was planning to invest a billion US\$ in electronic cigarettes [32]. The estimates of electronic cigarette sales are forecast to reach US\$419.6 million by 2022 [16].

The study revealed that electronic cigarette smoking was associated with the perception of electronic cigarettes as a smoking cessation aid. This may relate to the campaign from the electronic cigarettes industry that the device could support people to quit cigarette smoking dependence [33]. In 2017, the Indonesian FDA monitored the marketing activities of electronic cigarette sellers in online marketplaces, social media, video marketing and vape shops. The results found that most of the sellers used health claims, specifically claiming that they promoted electronic cigarettes as a quit smoking device, were a healthy alternative to using tobacco and were environmentally friendly [17]. This result suggests that adolescents were susceptible to promotional activities.

More than one-third of adolescents perceived that electronic cigarettes were easily found. Multivariate analysis revealed that school locations and availability were important predictors of ever having used electronic cigarettes. Electronic cigarette products are easily accessed through the online marketplace, social media and vape shops [17]. Downtown school locations may further ease access to vape stores. Based on the brief observation in Vapemagz’s websites, the vape store locations are more concentrated in city centers than in suburban areas [34]. The use of zoning to restrict the density and location of electronic cigarette retailers may be effective in reducing the availability of electronic cigarettes. Zoning laws have been widely used to restrict the operations of alcohol and fast food retailers [35,36]. A study in the US found that zoning laws successfully reduced alcohol availability and alcohol-related problems such as liver cirrhosis and motor vehicle crashes [35]. The government should consider how the land use restriction model could be extended to vape stores.

Predictors	AOR (95% CI)	p-value
Male sex	3.522 (2.316–5.357)	<0.001
Downtown school location	1.711 (1.153–2.538)	0.008
Lifetime cigarette smoker	8.740 (5.126–14.901)	<0.001
Current cigarette smoker	18.380 (10.577–31.938)	<0.001
Peer use	2.073 (1.311–3.277)	0.002
Availability	2.370 (1.526–3.679)	<0.001
Aid to quit cigarettes	5.197 (2.099–13.106)	<0.001

Table 4.
Predictors of ever
using electronic
cigarettes

The participants who had a peer using electronic cigarettes were more likely to use electronic cigarettes. The result is in line with prior studies in Poland, South Carolina, and Hong Kong [24,37,38]. The previous meta-analysis showed that the association with peer smoking was greater compared to family members [39]. A laboratory-based behavioral study found that the risky decisions of adolescents was influenced by the choices of their peers [40]. Furthermore, adolescents tended to imitate conventional cigarette smoking behaviors of others that were associated with their social context and shared a common normative belief [41]. It is noteworthy that the evidence of peer influence on tobacco product use behavior was extended to electronic cigarette use in this study.

Consistent with previous studies [42,43], we found that electronic cigarette use was associated with current conventional cigarette smoking status. There may be some reasons why combustible cigarette smokers take up electronic cigarettes. First, some electronic cigarettes' mold resembles traditional cigarettes [5]. This will ease current conventional cigarette smokers to experiment with electronic cigarettes. Second, most electronic cigarette industries promote the product as a smoking cessation aid or a healthier alternative. Thus, conventional smokers may exert electronic cigarettes to reduce the number of conventional cigarettes smoked. Third, electronic cigarettes are generally more accepted by the public than conventional cigarettes [44]. A prior study in a Swiss-Italian Middle school found that approval from other people was associated with tobacco use among adolescents [41].

This study has some limitations. First, it was based on a cross-sectional design, so we are not able to infer whether independent variables caused electronic cigarette use or otherwise. Second, private and vocational high school students were excluded from the study. The provincial education office also did not allow data collection from 12th-grade students. Hence, the findings may not be generalized to those groups. Finally, we used the multistage cluster sampling method and thus may have some cluster effect.

Conclusion

The research has some implications for public health research and policy in Indonesia. First, electronic cigarette use among males is increasing rapidly in a relatively short period from less than 1% in 2011 to 8.2% in 2015. Therefore, the government should extend the tobacco control measures such as introducing smoke-free area enforcements, imposing pictorial health warnings, tobacco advertisement and sponsorship bans and prohibit sales to minors on all types of tobacco products and not just conventional cigarettes. Second, it is worth noting that about 3% of participants who used electronic cigarettes had never smoked conventional cigarettes before. Therefore, further study is necessary to investigate the role of electronic cigarettes in inducing adolescents to take up conventional cigarettes. Third, we found school locations and availability were significantly associated with having ever used electronic cigarettes. Accessibility may induce electronic cigarette use among adolescents. Hence, we suggest that the government introduces zoning laws to restrict vape store locations near schools and residential areas. Ultimately, multivariate analysis revealed that affordability was not significantly associated with electronic cigarette use. Hereinafter, more than a quarter of participants perceived that electronic cigarette prices were affordable. These facts may suggest that the price of electronic cigarettes is relatively cheap in Indonesia. Thus, we recommend the government to charge maximum tax rates for electronic cigarette products.

Conflict of interest: There is no conflict of interest.

References

1. Besaratinia A, Tommasi S. Vaping: a growing global health concern. *EClinicalMedicine*. 2019 Dec; 17: 100208. doi: [10.1016/j.eclinm.2019.10.019](https://doi.org/10.1016/j.eclinm.2019.10.019).
2. World Health Organization [WHO]. WHO report on the global tobacco epidemic 2019. Geneva: WHO; 2019.
3. Azagba S, Manzione L, Shan L, King J. Trends in smoking behaviors among US adolescent cigarette smokers. *Pediatrics*. 2020 Mar; 145(3). doi: [10.1542/peds.2019-3047](https://doi.org/10.1542/peds.2019-3047).
4. Eastwood B, Dockrell MJ, Arnott D, Britton J, Cheeseman H, Jarvis MJ, McNeill A. Electronic cigarette use in young people in Great Britain 2013-2014. *Public Health*. 2015 Sep; 129(9): 1150-6. doi: [10.1016/j.puhe.2015.07.009](https://doi.org/10.1016/j.puhe.2015.07.009).
5. Kementerian Kesehatan Republik Indonesia (MOH-Kemenkes). Laporan Nasional Riskesdas 2018. Jakarta: Lembaga Penerbit Badan Penelitian dan Pengembangan Kesehatan (LPB); 2019. [text in Indonesian].
6. Noel JK, Rees VW, Connolly GN. Electronic cigarettes: a new 'tobacco' industry? *Tob Control*. 2011 Jan; 20(1): 81. doi: [10.1136/tc.2010.038562](https://doi.org/10.1136/tc.2010.038562).
7. van der Tempel J, Noormohamed A, Schwartz R, Norman C, Malas M, Zawertailo L. Vape, quit, tweet? Electronic cigarettes and smoking cessation on Twitter. *Int J Public Health*. 2016 Mar; 61(2): 249-56. doi: [10.1007/s00038-016-0791-2](https://doi.org/10.1007/s00038-016-0791-2).
8. Kalkhoran S, Glantz SA. E-cigarettes and smoking cessation in real-world and clinical settings: a systematic review and meta-analysis. *Lancet Respir Med*. 2016 Feb; 4(2): 116-28. doi: [10.1016/S2213-2600\(15\)00521-4](https://doi.org/10.1016/S2213-2600(15)00521-4).
9. Kaiser MA, Prasad S, Liles T, Cucullo L. A decade of e-cigarettes: limited research & unresolved safety concerns. *Toxicology*. 2016 Jul; 365: 67-75. doi: [10.1016/j.tox.2016.07.020](https://doi.org/10.1016/j.tox.2016.07.020).
10. Tzortzi A, Kapetanstrataki M, Evangelopoulou V, Beghrakis P. A systematic literature review of e-cigarette-related illness and injury: not just for the respirologist. *Int J Environ Res Public Health*. 2020 Mar; 17(7). doi: [10.3390/ijerph17072248](https://doi.org/10.3390/ijerph17072248).
11. U.S. Department of Health and Human Services. E-cigarette use among youth and young adults: areport of the Surgeon General. Atlanta, GA: U.S. Department of Health and Human Services; 2016.
12. Pisinger C, Dossing M. A systematic review of health effects of electronic cigarettes. *Prev Med*. 2014 Dec; 69: 248-60. doi: [10.1016/j.ypmed.2014.10.009](https://doi.org/10.1016/j.ypmed.2014.10.009).
13. Callahan-Lyon P. Electronic cigarettes: human health effects. *Tob Control*. 2014 May; 23(Suppl 2): ii36-40. doi: [10.1136/tobaccocontrol-2013-051470](https://doi.org/10.1136/tobaccocontrol-2013-051470).
14. Lestari KS, Humairo MV, Agustina U. Formaldehyde vapor concentration in electronic cigarettes and health complaints of electronic cigarettes smokers in Indonesia. *J Environ Public Health*. 2018; 2018: 9013430. doi: [10.1155/2018/9013430](https://doi.org/10.1155/2018/9013430).
15. Fernandez E, Ballbe M, Sureda X, Fu M, Salto E, Martinez-Sanchez JM. Particulate matter from electronic cigarettes and conventional cigarettes: a systematic review and observational study. *Curr Environ Health Rep*. 2015 Dec; 2(4): 423-9. doi: [10.1007/s40572-015-0072-x](https://doi.org/10.1007/s40572-015-0072-x).
16. Orlan EN, Parascandola M, Grana R. JUUL from the USA to Indonesia: implications for expansion to LMICs. *Tob Control*. 2019 Jul. doi: [10.1136/tobaccocontrol-2019-054979](https://doi.org/10.1136/tobaccocontrol-2019-054979) (article in press).
17. Badan Pengawas Obat dan Makanan. Kajian rokok elektronik di Indonesia. Jakarta: Direktorat Napza BPOM; 2017 [text in Indonesian].
18. Laestadius LI, Wahl MM, Pokhrel P, Cho YI. From Apple to Werewolf: a content analysis of marketing for e-liquids on Instagram. *Addict Behav*. 2019 Apr; 91: 119-27. doi: [10.1016/j.addbeh.2018.09.008](https://doi.org/10.1016/j.addbeh.2018.09.008).
19. Huang J, Tauras J, Chaloupka FJ. The impact of price and tobacco control policies on the demand for electronic nicotine delivery systems. *Tob Control*. 2014 Jul; 23(Suppl 3): iii41-7. doi: [10.1136/tobaccocontrol-2013-051515](https://doi.org/10.1136/tobaccocontrol-2013-051515).

20. Biliblicom. 2020. [cited 2020 April 13]. Available at: <https://www.bibli.com/jual/rokok-elektrik?page=1&start=0&searchTerm=rokok%20elektrik&intent=false&merchantSearch=true&multiCategory=true&customUrl=rokok-elektrik&sort=0&category=PE-1000540&catIntent=false>.
21. Lian TY, Dorotheo U. The tobacco control atlas: ASEAN region. 4th ed. Bangkok: Southeast Asia Tobacco Control Alliance (SEATCA); 2018.
22. Bigwanto M, Nurmansyah MI, Orlan E, Farradika Y, Purnama TB. Determinants of e-cigarette use among a sample of high school students in Jakarta, Indonesia. *Int J Adolesc Med Health*. 2019 Dec. doi: [10.1515/ijamh-2019-0172](https://doi.org/10.1515/ijamh-2019-0172) (article in press).
23. Data Referensi Pendidikan. 2014. [cited 2014 October 24] Available at: <http://referensi.data.kemdikbud.go.id/>.
24. Kaleta D, Wojtysiak P, Polanska K. Use of electronic cigarettes among secondary and high school students from a socially disadvantaged rural area in Poland. *BMC Public Health*. 2016 Aug; 15: 703. doi: [10.1186/s12889-016-3417-y](https://doi.org/10.1186/s12889-016-3417-y).
25. Hamilton HA, Ferrence R, Boak A, Schwartz R, Mann RE, O'Connor S, Adlaf EM. Ever use of nicotine and nonnicotine electronic cigarettes among high school students in Ontario, Canada. *Nicotine Tob Res*. 2015 Oct; 17(10): 1212-8. doi: [10.1093/ntr/ntu234](https://doi.org/10.1093/ntr/ntu234).
26. Lee S, Grana RA, Glantz SA. Electronic cigarette use among Korean adolescents: a cross-sectional study of market penetration, dual use, and relationship to quit attempts and former smoking. *J Adolesc Health*. 2014 Jun; 54(6): 684-90. doi: [10.1016/j.jadohealth.2013.11.003](https://doi.org/10.1016/j.jadohealth.2013.11.003).
27. Institute for Global Tobacco Control Johns Hopkins Bloomberg School of Public Health. Country laws regulating e-cigarettes: a policy scan. 2019. [cited 2019 December 18]. Available at: https://www.globaltobaccocontrol.org/e-cigarette_policyscan.
28. Palipudi KM, Mbulo L, Morton J, Mbulo L, Bunnell R, Blutcher-Nelson G, Kosen S, Tee GH, Abdalla AME, Al Mutawa KA, Barbouni A, Antoniadou E, Fouad H, Khoury RN, Rarick J, Shinha DN, Asma S, GATS Collaborative Group. Awareness and current use of electronic cigarettes in Indonesia, Malaysia, Qatar, and Greece: findings from 2011-2013 global adult tobacco surveys. *Nicotine Tob Res*. 2016 Apr; 18(4): 501-7. doi: [10.1093/ntr/ntv081](https://doi.org/10.1093/ntr/ntv081).
29. Hurt RD, Ebbert JO, Achadi A, Croghan IT. Roadmap to a tobacco epidemic: transnational tobacco companies invade Indonesia. *Tob Control*. 2012 May; 21(3): 306-12. doi: [10.1136/tc.2010.036814](https://doi.org/10.1136/tc.2010.036814).
30. BPS-Statistics Indonesia. Statistical yearbook of Indonesia 2019. Jakarta: BPS-Statistics Indonesia; 2020.
31. Ntoma RR. Sejarah vape; sebuah ide hingga jadi industri milyaran dollar. *Vapemagz Indonesia*. 2018 August. [cited 2020 April 22]. Available at: <https://vapemagz.co.id/wp-content/uploads/2018/08/Vapemagz-Agustus-2018.pdf>.
32. Khadafi M. Bertemu Jokowi, Phillip Morris lirik pasar rokok elektrik Indonesia. 2020. [updated 2020 February 28; cited 2020 April 22]. Available at: <https://market.bisnis.com/read/20200228/192/1207126/bertemu-jokowi-phillip-morris-lirik-pasar-rokok-elektrik-indonesia>.
33. Ramamurthi D, Gall PA, Ayoub N, Jackler RK. Leading-brand advertisement of quitting smoking benefits for e-cigarettes. *Am J Public Health*. 2016 Nov; 106(11): 2057-63. doi: [10.2105/AJPH.2016.303437](https://doi.org/10.2105/AJPH.2016.303437).
34. Vape Store Directory. *Vapemagz Indonesia*. 2020. [cited 2020 April 22] Available at: <https://vapemagz.co.id/search-result/?location=jakarta&category=1338&radius=25>.
35. Ashe M, Jernigan D, Kline R, Galaz R. Land use planning and the control of alcohol, tobacco, firearms, and fast food restaurants. *Am J Public Health*. 2003 Sep; 93(9): 1404-8. doi: [10.2105/ajph.93.9.1404](https://doi.org/10.2105/ajph.93.9.1404).
36. Lopez RP, Hynes HP. Obesity, physical activity, and the urban environment: public health research needs. *Environ. Health*. 2006 Sep; 5: 25. doi: [10.1186/1476-069X-5-25](https://doi.org/10.1186/1476-069X-5-25).

-
37. Barrington-Trimis JL, Berhane K, Unger JB, Cruz TB, Huh J, Leventhal AM, Urman R, Wang K, Howland S, Gilreath TD, Chou CP, Pentz MA, McConnell R. Psychosocial factors associated with adolescent electronic cigarette and cigarette use. *Pediatrics*. 2015 Aug; 136(2): 308-17. doi: [10.1542/peds.2015-0639](https://doi.org/10.1542/peds.2015-0639).
 38. Jiang N, Wang MP, Ho SY, Leung LT, Lam TH. Electronic cigarette use among adolescents: a cross-sectional study in Hong Kong. *BMC Public Health*. 2016 Mar; 16: 202. doi: [10.1186/s12889-016-2719-4](https://doi.org/10.1186/s12889-016-2719-4).
 39. Wang JW, Cao SS, Hu RY. Smoking by family members and friends and electronic-cigarette use in adolescence: a systematic review and meta-analysis. *Tob Induc Dis*. 2018; 16: 5. doi: [10.18332/tid/84864](https://doi.org/10.18332/tid/84864).
 40. Tomova L, Pessoa L. Information about peer choices shapes human risky decision-making. *Sci Rep*. 2018 Apr; 8(1): 5129. doi: [10.1038/s41598-018-23455-7](https://doi.org/10.1038/s41598-018-23455-7).
 41. Scalici F, Schulz PJ. Parents' and peers' normative influence on adolescents' smoking: results from a Swiss-Italian sample of middle schools students. *Subst Abuse Treat Prev Policy*. 2017 Jan; 12(1): 5. doi: [10.1186/s13011-017-0089-2](https://doi.org/10.1186/s13011-017-0089-2).
 42. Lee JA, Lee S, Cho HJ. The relation between frequency of e-cigarette use and frequency and intensity of cigarette smoking among South Korean adolescents. *Int J Environ Res Public Health*. 2017 Mar; 14(3): 305. doi: [10.3390/ijerph14030305](https://doi.org/10.3390/ijerph14030305).
 43. Nadasan V, Foley KL, Penzes M, Paulik E, Mihaicuta S, Abram Z, Balint J, Urban R. Use of electronic cigarettes and alternative tobacco products among Romanian adolescents. *Int J Public Health*. 2016 Mar; 61(2): 199-207. doi: [10.1007/s00038-015-0774-8](https://doi.org/10.1007/s00038-015-0774-8).
 44. Trumbo CW, Harper R. Use and perception of electronic cigarettes among college students. *J Am Coll Health*. 2013; 61(3): 149-55. doi: [10.1080/07448481.2013.776052](https://doi.org/10.1080/07448481.2013.776052).

Corresponding authorRidhwan Fauzi can be contacted at: ridhwanfauzi@umj.ac.id

For instructions on how to order reprints of this article, please visit our website:

www.emeraldgrouppublishing.com/licensing/reprints.htm

Or contact us for further details: permissions@emeraldinsight.com

Factors associated with electronic cigarettes use among adolescents in Jakarta, Indonesia

ORIGINALITY REPORT

19%

SIMILARITY INDEX

10%

INTERNET SOURCES

13%

PUBLICATIONS

10%

STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to Nexford Learning Solutions Student Paper	2%
2	www.thaiscience.info Internet Source	2%
3	link.springer.com Internet Source	2%
4	Submitted to Illinois Central College Student Paper	1%
5	Submitted to University of Melbourne Student Paper	1%
6	Submitted to Herzing University Student Paper	1%
7	www.ncbi.nlm.nih.gov Internet Source	1%
8	Miaw Yn Jane Ling, Ahmad Farid Nazmi Abdul Halim, Dzulfitee Ahmad, Norfazilah Ahmad, Nazarudin Safian, Azmawati Mohammed Nawi. "Prevalence and Associated Factors of E-Cigarette Use among Adolescents in Southeast Asia: A	1%

Systematic Review", International Journal of Environmental Research and Public Health, 2023

Publication

-
- | | | |
|---|--|-----|
| 9 | afri-can.org
Internet Source | 1 % |
|---|--|-----|
-
- | | | |
|----|---|-----|
| 10 | Chakkraphan Phetphum, Atchara Prajongjeep, Kanyarat Thawatchaijareonying, Thanchanok Wongwuttiyan et al. "Personal and perceptual factors associated with the use of electronic cigarettes among university students in northern Thailand", Tobacco Induced Diseases, 2021
Publication | 1 % |
|----|---|-----|
-
- | | | |
|----|---|------|
| 11 | Ridhwan Fauzi, Sathirakorn Pongpanich. "The effect of price on cigarette consumption among youth in Indonesia: Implications for tobacco tax policy", World Medical & Health Policy, 2022
Publication | <1 % |
|----|---|------|
-
- | | | |
|----|--|------|
| 12 | Jeff R. Temple, Ryan C. Shorey, Yu Lu, Elizabeth Torres, Gregory L. Stuart, Vi D. Le. "E-cigarette use of young adults motivations and associations with combustible cigarette alcohol, marijuana, and other illicit drugs", The American Journal on Addictions, 2017
Publication | <1 % |
|----|--|------|
-
- | | | |
|----|--|------|
| 13 | www.degruyter.com
Internet Source | <1 % |
|----|--|------|
-

14	www.drugsandalcohol.ie Internet Source	<1 %
15	bmcpublichealth.biomedcentral.com Internet Source	<1 %
16	Submitted to Massachusetts College of Pharmacy & Allied Health Sciences Student Paper	<1 %
17	www.jmir.org Internet Source	<1 %
18	Submitted to Oklahoma State University Student Paper	<1 %
19	Submitted to Surabaya University Student Paper	<1 %
20	Submitted to University of Queensland Student Paper	<1 %
21	bmchealthservres.biomedcentral.com Internet Source	<1 %
22	journals.sagepub.com Internet Source	<1 %
23	www.frontiersin.org Internet Source	<1 %
24	Submitted to Adtalem Global Education Student Paper	<1 %
25	Thamil Arasu Saminathan, Wan Shakira Rodzlan Hasani, Tania Gayle Robert Lourdes, Muhammad Fadhli Mohd Yusoff et	<1 %

al. "Cessation of E-Cigarette Use Among Adolescents and Its Associated Factors", *Asia Pacific Journal of Public Health*, 2019

Publication

26

Wang, Man Ping, William Ho Cheung Li, Nan Jiang, Lai Yan Chu, Antonio Kwong, Vienna Lai, and Tai Hing Lam. "E-Cigarette Awareness, Perceptions and Use among Community-Recruited Smokers in Hong Kong", *PLoS ONE*, 2015.

Publication

27

www.jpmpmh.org

Internet Source

<1 %

28

www.mdpi.com

Internet Source

<1 %

29

Jaana M. Kinnunen, Hanna Ollila, Jaana Minkkinen, Pirjo L. Lindfors, Arja H. Rimpelä. "A Longitudinal Study of Predictors for Adolescent Electronic Cigarette Experimentation and Comparison with Conventional Smoking", *International Journal of Environmental Research and Public Health*, 2018

Publication

<1 %

30

Jina Choo, Songwhi Noh, Jihyun Moon, Jinah Park, Yoonjoo Jeoung, Wonji Song. "Intention to quit electronic cigarette smoking among university students who are e-cigarette users", *Journal of American College Health*, 2022

<1 %

31 Heni Trisnowati, Hesti Yuningrum, Naomi Nisari Rosdewi, Enge Surabina Ketaren. "Electronic Cigarette Use Patterns And Its Determinants Among Adolescents In Yogyakarta City-Indonesia", Jurnal Formil (Forum Ilmiah) Kesmas Respati, 2022
Publication

32 Luoja Dai, Weiyi Lu, Juanjuan Wang, Lulu Zhang, Jingfen Zhu. "Social environment exposure to electronic cigarettes and its association with e-cigarette use among adolescents in Shanghai, China", Frontiers in Public Health, 2022
Publication

33 Meng Wang, Ru-Ying Hu, Jin Pan, Hao Wang, Min Yu, Kai-Xu Xie, Wei-Wei Gong. "Awareness, current use of electronic cigarettes and associated smoking factors in Zhejiang Chinese adolescents", PLOS ONE, 2019
Publication

34 S. F. L. Kirk. "Characterizing the obesogenic environment: the state of the evidence with directions for future research", Obesity Reviews, 02/2010
Publication

35 Valentin Nădășan, Kristie L. Foley, Melinda Péntzes, Edit Paulik et al. "Use of electronic cigarettes and alternative tobacco products

among Romanian adolescents",
International Journal of Public Health, 2016

Publication

36

Yuki Kuwabara, Aya Kinjo, Maya Fujii, Aya Imamoto et al. "Heat-Not-Burn Tobacco, Electronic Cigarettes, and Combustible Cigarette Use Among Japanese Adolescents: A Nationwide Population Survey 2017", Research Square Platform LLC, 2019

Publication

<1 %

37

Heba Jafar Sabbagh, Wafaa Abdelaziz, Maryam Quritum, Nada AbuBakr AlKhateeb et al. "Cigarettes' use and capabilities-opportunities-motivation-for-behavior model: a multi-country survey of adolescents and young adults", Frontiers in Public Health, 2022

Publication

<1 %

38

Mouhamad Bigwanto, Mochamad Iqbal Nurmansyah, Elizabeth Orlan, Yoli Farradika, Tri Bayu Purnama. "Determinants of e-cigarette use among a sample of high school students in Jakarta, Indonesia", International Journal of Adolescent Medicine and Health, 2022

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Factors associated with electronic cigarettes use among adolescents in Jakarta, Indonesia

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10
