

paper ice cream

by Susanty Susanty

Submission date: 05-Mar-2022 08:56PM (UTC+0700)

Submission ID: 1777114304

File name: Prosiding_Ice_Cream_FTUMJ_2021_-_Penulis_1.pdf (166.74K)

Word count: 2510

Character count: 12486

Determination of Used Cooking Oil's Toxicity Level by Adding Nutmeg Leaves (*Myristica Fragrans Houtt*) Extract

Susanty^{1*}, K. Chanda Dewi², Tri Yuni Hendrawati³, Wenny Diah Rusanti⁴, Fatma Sari⁵

^{1,2,3,4,5} Department of Chemical Engineering, Faculty of Engineering, Universitas Muhammadiyah Jakarta, Indonesia
Susanty@umj.ac.id

ABSTRACT

The nutmeg plant (*Myristica fragrans Houtt*) is one of the plants in the *Myristicaceae* family. The leaves of the nutmeg plant are rich in benefits, one of which has toxic indicating properties. The purpose of this study was to determine the level of used cooking oil toxicity by adding nutmeg leaf extract. This study used the maceration extraction method and the yield was 20%. Toxicity testing was carried out using the Brine Shrimp Lethality Test (BSLT) method. The research variable was the concentration of nutmeg leaf extract added into refined used cooking oil at 10 ppm, 50 ppm, 100 ppm, 150 ppm, 200 ppm, 250 ppm, and 300 ppm. Based on the research results, the optimum concentration of nutmeg leaf extract to determine the level of toxicity in used cooking oil was 250 ppm with an LC₅₀ value of 5.2389 ppm.

Keywords: Brine Shrimp Lethality Test (BSLT), nutmeg (*Myristica fragrans Houtt*) leaves, used cooking oil, toxicity.

© 2021 ICECREAM. All rights reserved

1. Introduction

Nutmeg (*Myristica fragrans Houtt*) is one of the plants in the *Myristicaceae* family. Due to their high value as a spice, nutmeg fruit and seeds have been an important trading commodity since Roman times. Nutmeg leaf is one part of the plant that has not been widely used. Compounds contained in nutmeg leaves include alkaloids, triterpenoids, tannins, and 2 flavonoids. The flavonoid components found in nutmeg leaves indicate the presence of phenolic compounds. This phenolic component is important considering its large role in the treatment and prevention of disease, including as antioxidants and has cytotoxic properties [1].

Cooking oil is one of the most vital household necessities and is increasing. The function of cooking oil is as an ingredient in food processing. Repeated use of cooking oil can cause bad effects on health. This is because cooking oil is damaged due to hydrolysis, oxidation, polymerization, and browning reactions when used for frying. The oxidation and polymerization processes damage some vitamins and essential fatty acids in the oil so that it can cause poisoning in the body and various diseases, such as diarrhea, deposition of fat in blood vessels and cancer [2].

One of the efforts that can be done is by reprocessing cooking oil that has been used repeatedly in order to minimize the potential harmful to the body. For example, the use of

leaves of the nutmeg plant (*Myristica fragrans Houtt*) as an indicator in testing the toxicity of used cooking oil. Usually the method of testing the toxicity properties of a material or compound is the Brine Shrimp Lethality Test (BSLT) method [3].

Previous research by Puspa, et al used Phytochemical and toxicity test of nutmeg essential oil (*Myristica fragrans Houtt*) from Kumutan Island with a comparison of ethyl acetate and ethanol solvent using the steam distillation and toxicity testing using the Brine Shrimp Lethality Test (BSLT) method. It was found that the quality of the nutmeg leaf essential oil obtained, namely the refractive index value achieved of 1.475 according to SNI 06-2388-2006 and the results of the toxicity test showed an LC₅₀ value of 5.192 ppm. Based on phytochemical test results, namely Nutmeg leaf essential oil contains terpenoids, flavonoids and saponins [4].

11

2. Material and Methods

2.1. Materials and tools

The materials were cooking oil, nutmeg leaf species *Myristica fragans Houtt*, magnesium, HCl, Tween 80, ethanol 96%, *Artemia salina Leach* larvae, aquades.

The tools were balance sheet, beaker glass, test tube, 1000 ml beaker glass, 25 ml measuring glass, 250 ml measuring glass, 100 ml volumetric flask, funnel, stirring rod, blender, filter, aluminum foil, filter paper, 50 ml beaker glass, evaporator.

2.2. Methods

2.2.1. Maceration

Nutmeg leaves obtained from Parung. The determination process of nutmeg leaves at LIPI, Bogor. Then dehydration process of nutmeg leaves was performed under the sun

for 3 days until the color turns brown. The dried leaves were blended until smooth and filtered until a fine powder of nutmeg leaves is obtained. Maceration extraction was 96% ethanol solvent with a ratio of 1: 10 [5]. Maceration extraction was carried out for 5 x 24 hours [6]. Nutmeg leaf extract was concentrated a rotary evaporator to obtain a thick extract [7, 8]. The viscous extract was tested qualitatively. Phytochemical tests for flavonoid compounds and toxicity tests using the Brine Shrimp Lethality Test (BSLT) method [9].

2.2.2. Phytochemical test

1 g of extract had put into a test tube and dissolved by 3 ml of 70% ethanol, and had been shaken to produce an orange solution. Heated on a water bath for ± 1 minute. Filtered to separate the filtrate and residue. The filtrate was added with 0.1 grams of Mg metal and produced an orange solution and there was a yellow precipitate. Then, 2 drops of HCl were added. The red color change in the ethanol layer indicates the presence of flavonoid compounds [10].

2.2.3. Brine Shrimp Lethality Test (BSLT)

200 ml solution of 1000 ppm of used cooking oil after being purified were prepared. 100 ml of 1000 ppm nutmeg leaf extract solution. Was then add Concentrations of the extract within the solution of 1000 ppm, varied respectively 300 ppm, 250 ppm, 200 ppm, 150 ppm, 100 ppm, 50 ppm, and 10 ppm. Each sample was poured into 20 ml beaker glass. The solution of 1000 ppm of used cooking oil was poured into a 20 ml beaker glass. 5 ml of extract solution were dropped into the used cooking oil sample, and then stirred until homogeneous. 10 *Artemia Salina L* shrimp larva were put into each beaker glass containing sample solution of varying concentrations of extract and a sample solution of used cooking oil and extract

mixture. The sample were observed for 1 x 24 hours, and the dead *Artemia Salina L* shrimp larvae were accounted. LC₅₀ [11].

2.3. Analytical Method

Yield of the extraction of nutmeg leaves were by calculated according to this formula :

$$\text{Yield} = \frac{\text{nutmeg leaf extract weight}}{\text{dry nutmeg leaf weight}} \times 100 \%$$

Analysis of the data of the Toxicity test were calculated using the LC₅₀ formula.

$$y = a + bx$$

$$\text{LC}_{50} = \text{arc log } x$$

description :

a = intercept; y = probit value; x = log¹⁰ concentration (ppm); b = slope

The probit value is an analysis that can estimate the effective dose by determining the concentration of mortality. The log concentration of extracts were compared, as well as the number of larvae mortality with the value of the probits depicted in an XY graph. The X axis showed the log concentration of nutmeg leaf extract and the amount of mortality, while the Y axis showed the probits value of each of these concentration logs. So that it was known the optimum conditions where the addition of nutmeg leaf extract works optimally.

3. Results and Discussions

3.1. Results

3.1.1. Nutmeg Leaf Yield

The thick extract of nutmeg leaves was made by the maceration extraction method with 96% ethanol as a solvent achieved a yield of 20%.

3.1.2. Phytochemical Test Results

In determining the presence of flavonoid compounds in the nutmeg leaf extract, a qualitative test was carried out. The sample was heated for ± 1-2 minutes, 0.1 grams of Mg to form a precipitate were added, then 2 drops of concentrated HCl were added too. A positive result was confirmed when a red color appeared. Based on the research conducted, nutmeg leaf extract contains flavonoid compounds was indicated by a red color change during the test.

3.1.3. Brine Shrimp Lethality Test (BSLT) Results

Toxicity testing has been done using the Brine Shrimp Lethality Test (BSLT) method. The sample consisted of purified used cooking oil, and thick extract of nutmeg leaves. The test was carried out using artificial seawater media which was added to 10 tail *A. salina shrimp* larvae and counted the number of *A. salina shrimp* larvae that died after being left for 1 x 24 hours during the test.

Table 1. The results of BSLT test for nutmeg leaf extract *Myristica fragans Houtt*

No	Nutmeg leaf extract concentration (ppm)	Mortality (tail)	Total (<i>A. Salina larva</i>) (tail)
1	10	2	10
2	50	4	10
3	100	4	10
4	150	6	10
5	200	7	10
6	250	8	10
7	300	8	10

Table 1 and 2 show the results of the toxicity test of used cooking oil with the addition of nutmeg leaf extract using the Brine Shrimp Lethality Test (BSLT) method.

Table 2. BSLT test results of cooking oil mixing and nutmeg leaf extract mixture

No	Cooking oil and nutmeg leaf extract mixture concentration (ppm)	Mortality (tail)	Total (<i>A. Salina</i> larva) (tail)
1	10	6	10
2	50	7	10
3	100	7	10
4	150	8	10
5	200	8	10
6	250	8	10
7	300	9	10

3.2. Discussion

This research was carried out by utilizing nutmeg plants which were more specific on nutmeg leaves that thrive in the Bogor area, West Java with the species *Myristica fragrans Houtt.* The reason for choosing nutmeg leaves as research raw materials was that in addition to nutmeg plants that are fertile in the area, nutmeg leaves contain flavonoids, saponins, terpenoids and are toxic.

The following is a graph of the probits versus mortality values of nutmeg leaf extract and used cooking oil added with *Myristica fragrans Houtt* nutmeg leaf extract.




Figure 1. Results of Brine Shrimp Lethality Test nutmeg leaf extract

Based on the graph above, the Brine Shrimp Lethality Test of nutmeg leaf extract at 10 ppm had a mortality of 2 larvas with a probit value of 4.16, 50 ppm had a mortality of 4 larvas with a probit value of 4.75, 100 ppm had a mortality of 4 larvas with a probit value of 4.75, 150 ppm had a mortality of 6 larvas with a probit value of 5.52, 200 ppm had a mortality of 7 larvas with a probit value of 5.52, 250 ppm had a mortality of 8 larvas with a probit value of 5.84, and 300 ppm had a mortality of 8 larvas with the probits value is 5.84.




Figure 2. Results of Brine Shrimp Lethality Test for used cooking oil and nutmeg leaf extract mixture

Based on the Fig. 2, the Brine Shrimp Lethality Test of used cooking oil and nutmeg leaf extract mixture at 10 ppm had a mortality of 6 larvas with a probit value of 5.52, 50 ppm had a mortality of 7 larvas with a probit value of 5.52, 100 ppm had a mortality of 7 larvas with probits value of 5.52, 150 ppm had a mortality of 8 larvas with a probits value of 5.84, 200 ppm had a mortality of 8 larvas with a probit value of 5.84, 250 ppm had a mortality of 8 larvas with a probits value of 5.84, and 300 ppm had a mortality of 9 tails with a probit value of 6.28.

Fig. 3 shows the LC₅₀ of nutmeg leaf extract and used cooking oil added with nutmeg leaf extract.




Figure 3. LC₅₀ of nutmeg leaf extract

Fig. 4 shows the LC₅₀ of used cooking oil and nutmeg leaf extract mixture.




Figure 4. LC₅₀ of used cooking oil and nutmeg leaf extract mixture

Based on the analysis of the BSLT test, the LC₅₀ value of nutmeg leaf extract (*Myristica fragrans Houtt*) was 74.098 ppm and the LC₅₀ value of used cooking oil added to nutmeg leaf extract (*Myristica fragrans Houtt*) was 5.2389 ppm. Thus, the addition of nutmeg leaf extract (*Myristica fragrans Houtt*) can measure the level of toxicity of the used cooking oil.

4. Conclusion

The yield of nutmeg leaf extract using 1:10 maceration extraction method using 96% ethanol solvent was obtained as much as 20%. The optimum condition for the addition of nutmeg leaf extract in the used cooking oil toxicity test was at a concentration of 250 ppm. The results of the BSLT test of used cooking oil added with nutmeg leaf extract obtained an LC₅₀ value of 5.2389 ppm. The lower the LC₅₀ value, made the mortality rate of larvae *Artemia s L* higher.

Acknowledgement

Thank you to the Center for Research and Community Service (LPPM) of the Universitas Muhammadiyah Jakarta for funding this research through the Internal Research grant scheme of the Universitas Muhammadiyah Jakarta.

References

- [1] Fawwaz, M., & Siti Nurdiansya A, M. B. *Potensi Daun Pala (Myristica fragrans Houtt) Sebagai Sumbang Fenolik*. 2017. Jurnal Fitofarmaka Indonesia, p. 212-214.
- [2] Rahayu, L. H., & Sari Purnavita, H. Y. *Potensi Sabut dan Tempurung Kelapa sebagai Adsorben untuk Meregenerasi Minyak Jelantah*. 2014. Jurnal Momentum, p. 1-7.
- [3] Musyirna rahmah Nst, S. H. *Aktivitas Sitotoksik Ekstrak Heksana, Diklorometana Dan Metanol Daun Keli Beling (Sericalyx crispus. L) Terhadap Artemia salina Leach*. 2011. Jurnal Ilmu Farmasi dan Farmasi Klinik, p. 71-74.
- [4] Puspa, O. E., Syahbanu, I., & W, M. A. *Uji Fitokimia dan Toksisitas Minyak Atsiri Daun Pala (Myristica fragrans Houtt) dari Pulau Lemukutan*. 2017. JKK, p. 1-6.

[5] Fauzana, D. L. *Perbandingan Metode Maserasi, Remaserasi, Perkolasi Dan Reperkolasi Terhadap Rendemen Ekstrak Temulawak (Curcuma xanthorrhiza Roxb.)*. 2010. Bogor: Institut Pertanian Bogor.

[6] Sitepu, J. S. *Pengaruh Variasi Metode Ekstraksi Secara Maserasi Dan Dengan Alat Soxhlet Terhadap Kandungan Kurkuminoid Dan Minyak Atsiri Dalam Ekstrak Etanolik Kunyit (Curcuma domestica Val.)*. 2017. Yogyakarta: Universitas Sanata Dharma .

[7] Nuramalia, W. *Evaporator Design Pada Pabrik Sorbitol Menggunakan Bahan Baku Dekstrosa Dengan Proses Hidrogenasi Katalitik*. 2019. Semarang: Universitas Negeri Semarang

[8] Supriatna, A. *Uji Performasi Dan Analisa teknik Alat Evaporator Vakum*. 2008. Bogor: Institut Pertanian Bogor.

[9] Susmayantia, W., Fachriyaha, E., & Kusrinia, D. *Isolasi, Identifikasi dan Uji Sitotoksik Senyawa Flavonoid dari Ekstrak Etil Asetat Daun Binahong (Anredera Cordifolia (Tenns) Stennis)*. 2021. Journal of Scientific and Applied Chemistry , 88-93.

[10] Devi, N. S. *Uji Fitokimia Pada Ekstrak Rimpang Temulawak (Curcuma Zanthorrhiza)*. 2016. p. 8-13.

[11] Ginting, B., Hadijah, S., & Helwati, H. *Isolation Of Antioxidant Compounds Of N-Hexane Extract Of Nutmeg (Myristica fragrans Houutt)Leaves*. 2017. Jurnal Kimia Mulawarman, p. 46-52.

paper ice cream

ORIGINALITY REPORT

10%

SIMILARITY INDEX

8%

INTERNET SOURCES

5%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1	jurnal.umj.ac.id Internet Source	2%
2	etheses.uin-malang.ac.id Internet Source	1%
3	dokumen.pub Internet Source	1%
4	www.science.gov Internet Source	1%
5	www.sumobrain.com Internet Source	1%
6	eudl.eu Internet Source	1%
7	Edrada-Ebel, RuAngelie, and Marcel Jaspars. "The 9th European Conference on Marine Natural Products", Marine Drugs, 2015. Publication	1%
8	Submitted to Lovely Professional University Student Paper	<1%

journal.uniga.ac.id

9

Internet Source

<1 %

10

repository.ub.ac.id

Internet Source

<1 %

11

Vida Elsyana, Maria Bintang, Bambang Pontjo Priosoeryanto. " Cytotoxicity and Antiproliferative Activity Assay of Clove Mistletoe ((L.) Miq.) Leaves Extracts ", Advances in Pharmacological Sciences, 2016

Publication

<1 %

12

www.scilit.net

Internet Source

<1 %

13

Md. Harun-Or-Rashid ., Nurul Karim ., M.A. Gafur ., Md. Golam Sadik . et al. "Isolation and Biological Activities of Chemical Constituents from the Stems of Ipomoea turpethum", Pakistan Journal of Biological Sciences, 2006

Publication

<1 %

14

Nurhamidah, H Nurdin, Y Manjang, A Dharma, Suryati. " Cytotoxic activity assay from leaves and fruit extracts of (Miq.) using brine shrimp lethality test method ", Journal of Physics: Conference Series, 2018

Publication

<1 %

15

www.ncbi.nlm.nih.gov

Internet Source

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On